

SUBTHRESHOLD ρ^0 PHOTOPRODUCTION ON ^2H , ^3He and ^{12}C ¹

G.J. LOLOS

Department of Physics, University of Regina, Regina, SK, S4S 0A2, Canada

Abstract

The ρ^0 has been photoproduced using tagged photon energies in the 700-1120 MeV region on ^2H , ^3He and ^{12}C by utilizing the Fermi momentum of the bound nucleons to produce the ρ^0 . These energies lie mostly below the production threshold on the free proton. Large mass modifications are indicated from the analysis, together with large polarizations of the produced ρ^0 mesons with helicity zero, unlike the case of coherent ρ^0 production on nuclei.

1 Introduction

The effects of hadron density and/or temperature on the mass and lifetimes of vector mesons have been the subject of numerous investigations, experimental as well as theoretical [1]. Under conditions of chiral symmetry restoration, the quark condensate vanishes, $\langle \bar{q}q \rangle \rightarrow 0$. The condensate is not, however, an observable but it is related to the meson mass, which is. Thus, the behaviour of vector meson masses under high hadron densities and/or temperatures has become a testbed of our understanding of hadron or quark dynamics in nuclear matter. A renormalization of vector meson masses in hot and dense matter is of significant importance to the pursuit of quark gluon plasma research, has relevance to the equation of state for nuclear matter in supernovae explosions, and could affect lower energy nuclear physics, as well. It has been argued that the mass modification of vector mesons should be measurable even at normal nuclear matter densities [2].

Even though a reduction of vector meson mass in hadronic matter is supported by most of the more recent theoretical models, the spectral shape (manifested in the width) is more of an open question. If there is substantial broadening as reported in [3], then extracting meaningful mass change information from ρ production and decay experiments becomes problematic, especially so

¹Proceedings of the International Workshop XXVIII on Gross Properties of Nuclei and Nuclear Excitations, Hirschegg, Austria, January 16-22, 2000, p. 203.

for massive nuclei. In this respect, production of either the ω in massive nuclei or the ρ from light nuclei may be a better solution. In any case, production of vector mesons, by whatever method, is not enough to probe medium modifications. The production of mesons must be accompanied by a substantial decay fraction within the nuclear volume, a fact which favours either low beam energies or restricted phase space in which the vector mesons are essentially at rest with respect to the recoiling nuclear system.

2 The TAGX Results from $A(\gamma_t, \pi^+\pi^-)X$ Reactions

2.1 The Subthreshold Energy Regime

Photo- or lepto-production of vector mesons from nuclei has been and is being explored at many facilities. Given the quantum numbers of the vector mesons and those of the photon (real or virtual), the primary interaction of the latter proceeds via the interaction of the former with the nucleons in nuclei, the well established VMD model of γ -N interactions. At incident photon energies of approximately 1.2 GeV and higher, the diffractive mechanism dominates, in which the photon fluctuates into a ρ^0 or ω , which then scatters off the nucleon field with sufficient four-momentum transfer to bring the meson on the mass shell. In the case of nuclei, the single γ -N amplitudes add coherently and large production cross sections result. Since the momentum transfer t is less than $0.1 \text{ GeV}^2/c^2$, coherent or diffractive cross sections are very forward peaked. As a result, they also conserve helicity and the reaction is characterized by a large impact parameter of approximately 10 fm or more.

While coherent photoproduction produces vector mesons prolifically, the probability of the vector mesons penetrating the nuclear interior and decaying within it is rather small. The observation, then, of medium modifications in such reactions is diminished by “geometry” (including trajectory of the produced meson with respect to the nucleus) and the large Lorentz boost of the meson with respect to the nucleus. This is particularly severe for the ω due to its long $c\tau$ compared to that of the ρ^0 . One is thus forced to depend on the large number of mesons produced to observe the small fraction that penetrates and decays within the nuclear volume, and which is perhaps overwhelmed in the invariant mass signature from the low tail of the unmodified mesons decaying freely. Where the fraction is larger, like in the ρ -meson case, the free ρ^0 width is also large, thus effectively masking the modified mass. In the case of the ω , the narrow width makes it easier to separate the modified from the unmodified masses, but the fraction of the former to the latter is very small.

An alternative is to produce the ρ^0 below the free $\gamma + N \rightarrow \rho^0$ nominal threshold of 1083 MeV. In this case, the reaction is characterized by large t and the Fermi four-momentum of the struck nucleon is utilized to bring the ρ^0 on the mass shell. A penalty one pays for such *subthreshold* ρ^0 production is the small cross section compared to coherent production. There are numerous advantages, however. First, the ρ^0 production vertex is forced to be within the nuclear volume (small impact parameter) by the interaction of the photon with a bound nucleon. The reaction then is quasi-free γ -N interaction in nature. Second, the ρ^0 is produced with small relative momentum and Lorentz boost with respect to the struck nucleon or the nucleus, thus maximizing the probability of decay within the hadronic volume. Third, the large opening angle (of 180° in the ρ^0 rest frame) is essentially preserved in the lab frame due to the small Lorentz boost and this provides a characteristic signature of the ρ^0 decay, which can be used effectively in the suppression of competing processes.

2.2 The TAGX Results on ^3He

The choice of ^3He was made based on several reasons. (a): Its core nuclear density is near saturation density for heavier nuclei, while the small number of nucleons reduces final state interactions (FSI) between the pions and the recoiling nucleus or nucleons. (b): The kinematical analysis is much simpler due to the small number of spectator nucleons and the Monte Carlo (MC) simulations carry more confidence. And, (c): extrapolations of cross sections for background processes on proton and ^2H are more accurate for ^3He than more complex nuclei. Thus, it was felt that any mass modification, even if small, could carry higher confidence due to the simplicity of the probe-target system.

The analysis was based on a detailed and kinematically constrained fit to several distributions simultaneously for all processes that can lead to a $\pi^+\pi^-$ final state. Details are published in [4], [5], and [6]. The results were consistent with a large mass modification for the ρ^0 , which was also found to depend on incident photon energy. At photon energies as low as 660 ± 40 MeV, the best fit was obtained for a $m_{\rho^0}^* = 490 \pm 40$ MeV/ c^2 , while for 840 ± 40 MeV the mass of the ρ^0 was $m_{\rho^0}^* = 640 \pm 40$ MeV/ c^2 . For the highest energy bin available to the experiment (which was centered around 1080 MeV), the data were shown to be insensitive to any ρ^0 mass modification. The conclusions survived a number of different analyses, and even the data sets for two of the analyses were different and independent of each other.

The analysis was based on the simultaneous fitting of a number of “key” variables, such as missing momentum (p_{miss}), missing mass (m_{miss}), invariant mass ($m_{\pi\pi}$), pion opening angle ($\theta_{\pi\pi}$), and emission angle of the di-pion system (θ_{IM}). All these are kinematical observables and follow phase space distributions, applicable to each individual reaction assumed. For the 80 MeV wide photon energy bins in the analysis, the assumption was made that the matrix element is constant. The yield then is the product of several factors:

$$Y = |M^*(s, t)|^2 \cdot \Phi_{3He}(p) \cdot \Psi(s, I, J) \cdot [MBPS * F(m, \Gamma)] \quad (1)$$

where Y = Yield, $|M^*|$ = invariant matrix element for the reaction, Φ = the single nucleon momentum distribution, Ψ = function of spin, isospin, and total angular momentum, and $[MBPS * F]$ = the multi-body phase space for this reaction with the mass and width of any resonances folded in. Therefore, even though a variation of $|M^*|$ within the 80 MeV bin is not taken explicitly into account, the width Γ of the states involved has a large part of this energy dependence folded in. Each background and foreground reaction was simulated using Monte Carlo (MC) generators, which incorporated equation 1.

Cross sections for known processes, such as $\Delta\pi$, $N^*\pi$, $\Delta\Delta$ and $\pi^+\pi^-\pi^0$, were extracted and compared to established measurements in the literature, if such existed. The cross sections for ρ^0 production with modified masses were quite small (a few μb), while that of competing processes are of the order of 200 μb [5]. The small ρ^0 signal could be extracted due to the features of the TAGX acceptance. The trigger requirement of left-right (with respect to the photon beam) $\pi^+ - \pi^-$ coincidences, and the small out of plane acceptance of the spectrometer, result in substantially larger acceptance for the $\rho^0 \rightarrow \pi^+\pi^-$ decay than the other competing two-step processes leading to two-pion emission. Furthermore, the requirement of simultaneous fitting of so many different kinematical variables provided severe constraints to the strength of these background reactions in different regions of the various distributions.

One of the difficulties of the above analysis is the model dependency and the smallness of the ρ^0 production cross sections compared to all the other competing processes. The analysis, however, has been shown to be insensitive to reasonable variations in assumed cross sections for the dominant background processes because the ρ^0 events occupy quite distinct regions of phase space. Thus, variations in $\Delta\pi$ production cross sections, for example, affected $\Delta\Delta$ fits and resulting cross sections, but they did not affect the value of $m_{\rho^0}^*$. Another conclusion was that $m_{\pi\pi}$ is not a very sensitive parameter and good fits can be obtained for large differences in the input parameters. On the other hand, $\theta_{\pi\pi}$ and p_{miss} are difficult (and sensitive) distributions to fit.

In the case of coherent ρ^0 production and at very small angles of emission in the helicity frame of reference, the angular distribution for θ_π^* of one pion with respect to the di-pion center of mass momentum in the helicity frame, has a $\sin \theta_\pi^*$ -like distribution for unpolarized photons. This is a result of s -channel helicity conservation. In the case of helicity non-conservation, for example at large angles of ρ^0 emission due to large t exchange, a random distribution of polarization would result in an isotropic distribution for θ_π^* . When, however, such a $\theta_{\pi^+}^*$ distribution was formed, an enhancement in the regions of 0° and 180° was observed in the “raw” event sample. This has been interpreted as a substantial polarization component in the ρ^0 event sample which corresponds to the $l = 1, m = 0$ substate. Such a Y_1^0 -like distribution was most prominent in the same $m_{\pi\pi}$ bins where the independent kinematical analysis placed the mean of the modified ρ^0 mass.

2.3 The TAGX Results on ^2H , ^{12}C

The analysis of the ^2H and ^{12}C results is still ongoing and a final value for the modified masses has not been extracted yet. The philosophy and methodology is quite different than the analysis of the ^3He data. The simpler ^2H nucleus, coupled with photoabsorption cross sections in the literature for most of the channels leading to di-pion emission, allow the use of experimental amplitudes and cross sections and a likelihood method of fitting the kinematical observables. Unlike the analysis of the ^3He data, in this analysis the $\theta_{\pi^+}^*$ distributions are explicitly involved among the kinematical variables fitted. The cross section results are not yet final, however, the analysis based on the $\theta_{\pi^+}^*$ distribution is quite interesting. In Figure 1, the $\cos \theta_{\pi^+}^*$ distributions are shown for $^{12}\text{C} + ^2\text{H}$ data, (a) without any restrictions on the data, and (b) with a $\theta_{\pi\pi} \geq 120^\circ$ cut. Several observations can be drawn:

- In the $M_{\pi\pi}$ bin of 300-400 MeV/ c^2 , the shape of the distribution and the small number of events near the -1 and +1 limits is characteristic of acceptance effects due to trigger and geometry of TAGX. As such, the population in these two regions for the invariant mass bins in the 500-600 MeV/ c^2 regions, signify substantial $\rho^0 \rightarrow \pi^+\pi^-$ -like production.
- The application of the opening angle ($\theta_{\pi\pi}$) cut, removes events from the central regions of the distribution, but leaves the population around the two limits unchanged.

The $\cos \theta_{\pi^+}^*$ distributions for all three nuclei exhibit the same overall behaviour. The preliminary results from the cross section fitting of the ^2H data




Figure 1: Selected invariant mass bins for data from the ${}^2\text{H}, {}^{12}\text{C}(\gamma, \pi^+\pi^-)\text{X}$ experiment obtained with TAGX. The top panels have no $\theta_{\pi\pi}$ cut, whereas the bottom panels have been subjected to a $\theta_{\pi\pi} \geq 120^\circ$ cut. The invariant mass bins for 400-500 and 600-700 MeV/c^2 also exhibit a p -wave signature, but it's neither as strong nor as symmetric as that in the 500-600 MeV/c^2 bin; they are not shown here for the sake of clarity.

are consistent with a few μb cross section for polarized ρ^0 production in the $l = 1$, $m = 0$ substate and no strength in the $l = 1$, $m = 1$ substate. This is qualitatively in agreement with the ${}^3\text{He}$ data which also showed no statistically significant population in the small angle regions in plots such as shown in Figure 1. This was verified by successively applying increasingly tighter opening angle cuts and comparing the behaviour as a function of $\cos \theta_{\pi+}^*$ values.

2.4 The Effects of $\theta_{\pi\pi}$ Cuts

The $l = 1$, $m = 0$ signature is model independent and difficult to argue away. Nevertheless, there have been a number of possible explanations, other than a modified and polarized ρ^0 . One is that *there may be* other reactions which lead to two-pion emission in a relative p -state and with a $m = 0$ substate. However, no such specific reaction has ever been identified or proposed as




Figure 2: Deuterium distributions of the cosine of the pion angle in the helicity frame. The columns indicate the effect of progressively larger opening angle cut. The rows portray the different MC-generated channels, with the verticle arrow indicating that the particle in question is produced in a polarized state.

a candidate; in any case, if there is indeed one, this by itself is a new and interesting result. Another concern put forward is that the application of ever tighter $\theta_{\pi\pi}$ requirements may artificially produce such $l = 1$, $m = 0$ signatures. In order to address this valid concern, a detailed investigation of the behaviour of two of the most prolific background processes has been made as a function of $\theta_{\pi\pi}$ cuts. MC simulations of three production reactions have been made and analyzed in the same fashion as the real data are analyzed in the TAGX detector. These are shown in Figure 2 for ^2H , plotted as a function of $\cos\theta_H$ in the helicity frame.

The $\Delta\pi$ and $\Delta\Delta$ background channels are the two strongest channels leading to $\pi^+\pi^-$ production. As can be seen in Figure 2, whether the Δ is produced

polarized (indicated by the arrow) or not, the application of successively more demanding $\theta_{\pi\pi}$ cuts does not alter the shape of the distributions. In the case of unpolarized ρ^0 production, the distribution is essentially flat with a slight enhancement at the limits due to detector acceptance. Only in the case of polarized ρ^0 production do the MC simulations in Figure 2 agree with the observed behaviour of the data in Figure 1.

3 Conclusions

The analyses of ^3He data using MC and model dependent, as well as model independent variables, agree on a substantial mass modification of the ρ^0 . The preliminary results on ^2H and ^{12}C , using primarily the model independent $\theta_{\pi^+}^*$ analysis, agree with the earlier results and strongly support the production of longitudinally polarized (zero helicity) and mass modified ρ^0 -mesons. The similarities among the results for these three different nuclei is suggestive of a ρ^0 -N based interaction effect rather than a nuclear effect. Since the relative ρ^0 -N momenta at such low incident photon energies are also very low, the mean decay length of the ρ^0 from its production vertex is less than the nucleon radius. Hadronic densities at the *interior* of a nucleon are very high, thus qualitatively accounting for the large mass modifications reported so far from TAGX results. The excellent agreement on the $\theta_{\pi\pi}$ dependence for data and MC simulations conclusively shows that only a $l = 1$, $m = 0$ substate of the ρ^0 is present in the data. Such a production can be accounted for by a spin-flip transition for the target nucleon, consistent with a N^* excitation in the ρ^0 production process.

References

- [1] C.M. Ko *et al.*, Ann. Rev. Nucl. Sci. **47** (1997) 505.
- [2] G.E. Brown and M. Rho, Phys. Rev. Lett. **66** (1991) 2720.
- [3] F. Klinglet *et al.*, Nucl. Phys. **A560** (1997) 527.
- [4] G.J. Lolos *et al.*, Phys. Rev. Lett. **80** (1998) 241.
- [5] G.M. Huber *et al.*, Phys. Rev. Lett. **80** (1998) 5285.
- [6] M.A. Kagarlis *et al.*, Phys. Rev. C **60** (1999) 025203.